

Nursing

*ABCGN-CGRN
Certified Gastroenterology Registered Nurse*



For More Information – Visit link below:

<https://www.examsboost.com/>

Product Version

- ✓ **Up to Date products, reliable and verified.**
- ✓ **Questions and Answers in PDF Format.**

Latest Version: 6.0

Question: 1

Which of the following dairy products is MOST LIKELY to contain the least lactose for those who are lactose intolerant?

- A. Whole milk
- B. Cheese
- C. Regular whole-fat yogurt
- D. Greek-style whole-fat yogurt

Answer: B

Question: 2

Unless otherwise specified by manufacturer, multi-use vials that have been accessed and used should be discarded within which of the following time periods?

- A. One week
- B. 14 days
- C. 28days
- D. 60 days

Answer: B

Question: 3

Which of the following is a reversal agent for excessive sedation of a patient who has received a benzodiazepine?

- A. Atropine
- B. Romazicon (Flumazenil®)
- C. Naloxone (Narcan®)
- D. N-acetylcysteine

Answer: B

Question: 4

Following an esophagoscopy to obtain a biopsy of the thoracic esophagus, which of the following symptoms MOST indicates the need for emergent care for perforation?

- A. Chest pain, dysphagia, and tachycardia
- B. Mild cough and sore throat
- C. Nausea and vomiting
- D. Local discomfort but no systemic response

Answer: A

Question: 5

A 76-year-old female ate E. coli (O157:H7) contaminated vegetables and developed abdominal cramps and non-bloody diarrhea for 48 hours after which the diarrhea became bloody for 4 days. The patient is MOST at risk for developing which of the following?

- A. Intestinal necrosis
- B. Small bowel obstruction
- C. Intestinal perforation
- D. Hemolytic uremic syndrome

Answer: D

Question: 6

A patient taking metoclopramide has been prescribed haloperidol. For which of the following does this drug combination put the patient at increased risk?

- A. Tachycardia
- B. Tardive dyskinesia
- C. Excessive sedation
- D. Gbleeding

Answer: B

Question: 7

If a patient scheduled for a colonoscopy has a nose stud and enclosed lip ring, which of the following actions is appropriate?

- A. Remove the lip ring and tape the nose stud securely

- B. Leave both in place
- C. Remove both prior to the procedure
- D. Remove the nose stud but leave the lip ring in place

Answer: C

Question: 8

Cancer of the colon and rectum is primarily which of the following types of cancer?

- A. Adenocarcinoma
- B. Sarcoma
- C. Lymphoma
- D. Melanoma

Answer: A

Question: 9

According to the WHO three-step ladder approach to pain management, if a patient's abdominal pain associated with colon cancer varies from 4 to 8 on the pain scale, at which of the following steps should pain control be initiated?

- A. Step1
- B. Step2
- C. Step 3
- D. Whichever step is appropriate at the time of initiation

Answer: B

Question: 10

How soon after collection should duodenal aspirate be transported to the laboratory?

- A. Immediately
- B. Within 2 hours
- C. Within 4 hours
- D. Within 24 hours

Answer: A

Thank You for Trying Our Product

For More Information – **Visit link below:**

<https://www.examsboost.com/>

15 USD Discount Coupon Code:

G74JA8UF

FEATURES

- ✓ **90 Days Free Updates**
- ✓ **Money Back Pass Guarantee**
- ✓ **Instant Download or Email Attachment**
- ✓ **24/7 Live Chat Support**
- ✓ **PDF file could be used at any Platform**
- ✓ **50,000 Happy Customer**



Visit us at <https://www.examsboost.com/test/abcdn-cgrn/>