

Boost up Your Certification Score

Healthcare

ACMA-ACM

ACMA Accredited Case Manager (ACM)



For More Information – Visit link below:

<https://www.examsboost.com/>

Product Version

- ✓ **Up to Date products, reliable and verified.**
- ✓ **Questions and Answers in PDF Format.**

Visit us at: <https://www.examsboost.com/test/acma-acm>

Latest Version: 6.0

Question: 1

The case manager is investigating a case where a patient was restrained. The patient has filed a complaint, saying she was unlawfully restrained. The case management team was asked to investigate. The case manager has found:

- The patient was physically restrained with restraints
- The patient was physically harming himself and others

Which of the following is true?

- A. The patient was illegally restrained.
- B. The patient was legally restrained.
- C. The patient restrained himself.
- D. The patient asked for restraints.

Answer: B

Explanation:

Here's the expanded explanation: When considering the legality of using restraints on a patient, it is crucial to understand the context and the circumstances under which the restraints were applied. In this case, the key piece of information provided is that the patient was "physically harming himself and others." This behavior represents a clear and immediate risk to the safety of the patient and those around him, which justifies the use of restraints as a necessary and immediate response.

The use of restraints in healthcare settings is heavily regulated to protect patients' rights and ensure ethical treatment. Generally, restraints should only be used when absolutely necessary to prevent harm to the patient or others, and when less restrictive interventions have been ineffective. Given the scenario where the patient was actively harming himself and others, the application of restraints aligns with these guidelines.

Furthermore, the law and medical ethics require that the use of restraints must be followed by proper assessment, monitoring, and ongoing re-evaluation to ensure the continued need for restraint and to safeguard the patient's health and well-being. While the question does not specify whether these procedures were followed, the legality of the initial action of restraining the patient under these circumstances stands.

It's also important to note that patients do have the right to be free from restraints unless it is medically necessary. Here, the necessity arises from the immediate risk posed by the patient's actions, which legitimizes the use of restraints according to healthcare regulations and standards.

In conclusion, based on the information provided — specifically, that the patient was harming himself and others — the use of restraints was legally justified. However, any ongoing use of restraints would need to be regularly evaluated to ensure it remains justified and that the patient's rights and safety are upheld.

Question: 2

Motivational interviewing would include which of the following?

- A. Listening
- B. Implementing
- C. Evaluating
- D. Assessing

Answer: A

Explanation:

Motivational interviewing (MI) is a counseling approach developed in part by clinical psychologists William R. Miller and Stephen Rollnick. It is a directive, client-centered counseling style for eliciting behavior change by helping clients to explore and resolve ambivalence. Compared with non-directive counseling, it's more focused and goal-directed. The examination and resolution of ambivalence is a central purpose, and the counselor is intentionally directive in pursuing this goal.

MI is most commonly used to address addiction and the management of physical health conditions such as diabetes, heart disease, and asthma. This method particularly helps in motivating individuals to change certain behaviors that are not conducive to health or personal well-being.

Core to motivational interviewing is the concept of engaging clients in a collaborative relationship. Here are several key components that are typically included in motivational interviewing: 1. **Listening:**

This is a fundamental aspect of motivational interviewing. Effective listening in MI requires a deep, empathetic, and reflective listening style. The practitioner listens carefully to every word the client says, seeks to understand the client's perspective, and occasionally reflects back to the client what they have heard to ensure understanding and to provide the client with an opportunity to listen and reflect on their own words and feelings. 2. **Talking:** Dialogue in MI is client-centered. The practitioner talks less and encourages the client to express their thoughts and feelings. When practitioners do talk, they use open-ended questions that encourage discussion, reflection, and detailed responses from clients. This helps in exploring the reasons behind the clients' ambivalence towards behavior change. 3.

Implementing: Implementation refers to the stage where clients begin to take steps towards change. In motivational interviewing, practitioners support and facilitate this process by helping clients to set practical, achievable goals. This might involve discussing strategies for change, helping to plan for potential obstacles, and ensuring that the client feels capable of making the changes. 4. **Evaluating:** This involves reviewing the progress towards the goals. The practitioner helps the client to assess their successes and areas needing improvement. Evaluation is important as it provides both the client and practitioner with insights into what strategies are working and what adjustments might be needed to continue forward. 5. **Assessing:** Assessment in motivational interviewing involves continuously gauging the client's readiness and willingness to change. Practitioners use this process to tailor their approach to fit the client's current motivational levels, which can fluctuate. This might involve revisiting the benefits and costs of change as perceived by the client or exploring new sources of motivation.

In summary, motivational interviewing is a nuanced, dynamic, and client-centered approach that effectively supports individuals in making meaningful changes through a structured yet empathetic dialogue and collaborative relationship. It relies heavily on listening and talking but also systematically includes assessing, planning, implementing, and evaluating steps towards achieving behavioral change.

Question: 3

Jacob has come to the ER and was assessed. The physician found that Jacob has symptoms of shortness of breath, sharp chest pain, rapid heart rate, rapid breathing, and excessive sweating. A care plan has been planned out based on a diagnosis of a lung cancer. This care plan is not correct for Jacob. The case management team was asked to investigate why the plan was incorrectly done. The team found:

- The plan has incorrect actions for this patient
- The patient was misdiagnosed
- The patient was not assessed thoroughly

Given the information, what should the care plan be based on?

- A. Dementia.
- B. Pulmonary embolism.
- C. Throat cancer.
- D. Pneumonia.

Answer: B

Explanation:

The correct care plan for Jacob should be based on a diagnosis of pulmonary embolism, not lung cancer. This decision is reached after the case management team investigated the initial misdiagnosis and identified several issues in the initial assessment and care planning process. **Misdiagnosis:** Jacob's symptoms of shortness of breath, sharp chest pain, rapid heart rate, rapid breathing, and excessive sweating are more indicative of a pulmonary embolism than lung cancer. Pulmonary embolism is a critical condition caused by a blockage in one of the pulmonary arteries in the lungs, typically due to blood clots that travel to the lungs from the legs or other parts of the body. These symptoms are acute and align closely with the clinical presentation of a pulmonary embolism. **Incorrect Care Plan:** The initial care plan was designed for lung cancer, which typically includes treatments like surgery, chemotherapy, or radiation therapy, depending on the stage and type of cancer. However, these treatments are not appropriate for managing a pulmonary embolism. The care plan for pulmonary embolism should focus on immediately stabilizing the patient's condition, managing pain, and most importantly, dissolving or removing the clot. This typically involves the administration of anticoagulants (blood thinners), thrombolytic therapy (to break up clots), and in severe cases, surgical intervention to remove the clot. **Lack of Thorough Assessment:** It appears that the initial assessment did not include adequate diagnostic testing that could have revealed the presence of a pulmonary embolism. Tests such as a CT pulmonary angiography, a D-dimer test, or a ventilation-perfusion (V/Q) scan are crucial in diagnosing pulmonary embolism. The failure to perform these tests likely contributed to the incorrect diagnosis and inappropriate care plan. **Conclusion:** The correct diagnosis of pulmonary embolism is crucial because it is a potentially life-threatening condition that requires immediate and specific intervention. The care plan should be urgently revised to focus on anticoagulation therapy, monitoring for stability, and possibly procedures to remove the clot if indicated. This approach is essential to prevent further complications, including further embolic events or pulmonary hypertension, and to stabilize Jacob's condition effectively.

Question: 4

A case manager is examining continuum of care. RMNCH continuum of care generally ends at:

- A. Delivery.

- B. Pre-pregnancy.
- C. Childhood.
- D. Post pregnancy.

Answer: C

Explanation:

The RMNCH continuum of care is a comprehensive care approach that addresses the health needs of women, newborns, and children. It encompasses pre-pregnancy, pregnancy, childbirth, and the postnatal period, extending into childhood. This care model is designed to ensure a seamless transition across different stages of life and health care needs, promoting long-term well-being and survival.

The question specifically asks at which stage the RMNCH continuum of care generally ends.

Understanding the framework of RMNCH is crucial to answering this question. The RMNCH continuum starts from pre-pregnancy, which includes providing care and guidance before conception. This phase focuses on optimizing the health of women of reproductive age, preparing them for a healthy pregnancy. The next phase is pregnancy, where care is provided to maintain the health of the mother and the developing fetus.

Following pregnancy, the continuum includes childbirth, which involves managing labor and delivery to ensure the safety of both mother and newborn. The postnatal phase follows, where both the mother and the newborn receive care to promote recovery and healthy development. After the postnatal period, the continuum extends into childhood, addressing the health needs of the child as they grow.

Given this comprehensive approach, the RMNCH continuum of care generally ends at childhood. This is because the continuum is explicitly designed to provide integrated care from before pregnancy through to the child's development, ensuring a healthy start in life. After childhood, general health services continue to address the individual's needs, but they fall outside the specific RMNCH framework. Thus, the correct answer to the question would be "Childhood," as this marks the final phase of the specifically targeted RMNCH care continuum.

Question: 5

Checking for a disease when there are no symptoms is known as which of the following?

- A. screening
- B. laboratory testing
- C. sensitivity assessment
- D. none of the above

Answer: A

Explanation:

Screening refers to the process of checking for a disease or condition in individuals who do not exhibit any symptoms. This proactive health measure is crucial because it helps in identifying diseases early in their course before any clinical signs become apparent. By detecting a condition early, screening provides a chance for interventions that can prevent the disease from developing further or reduce its impact significantly.

Screening is typically conducted on populations that are considered at risk for certain diseases but who are currently asymptomatic. This can include screening for high cholesterol, certain cancers (such as breast cancer or colorectal cancer), diabetes, hypertension, and more. The methods used for screening vary depending on the disease but often include blood tests, imaging (like mammograms or colonoscopies), and other diagnostic procedures.

It is important to distinguish screening from diagnostic tests, which are performed on individuals who already show symptoms of a disease. Screening aims at primary or secondary prevention and is usually applied to a broader, asymptomatic population. Primary prevention aims to prevent the disease from occurring, while secondary prevention focuses on early detection and treatment, thereby stopping the progression of the disease.

The effectiveness of a screening program depends on several factors, including the sensitivity and specificity of the screening test, the prevalence of the disease in the population being screened, and the availability of effective treatment after detection. Sensitivity refers to the test's ability to correctly identify those with the disease, whereas specificity refers to the test's ability to correctly identify those without the disease.

Ultimately, the goal of screening is to improve public health outcomes by reducing the burden of disease and preventing avoidable complications and mortalities. This makes screening a key component of preventive medicine and public health strategies. It is essential, however, for screening programs to be carefully planned and implemented, considering both the benefits and potential harms, such as false positives, overdiagnosis, and the anxiety associated with testing.

Question: 6

Under the Health Insurance Portability and Accountability Act (HIPAA) any individual, organization or facility which meets the definition of “covered entity” must do certain things. Which of the following may NOT be a covered entity under HIPAA?

- A. free health clinic
- B. healthcare clearinghouses
- C. health plans
- D. healthcare providers who transmit health information in electronic form in connection with a transaction

Answer: A

Explanation:

Under the Health Insurance Portability and Accountability Act (HIPAA), a "covered entity" refers to any individual, organization, or facility that must comply with HIPAA's rules. These entities typically handle protected health information (PHI) and are involved in healthcare transactions that require privacy and security protections. The categories of covered entities under HIPAA include health plans, healthcare clearinghouses, and healthcare providers who conduct certain financial and administrative transactions electronically.

Among the choices given—free health clinic, healthcare clearinghouses, health plans, and healthcare providers who transmit health information in electronic form—the one that may not necessarily be a covered entity is the free health clinic. The primary reason for this is the nature of billing and transactions conducted by the entity.

Healthcare clearinghouses process nonstandard health information they receive from another entity into a standard format or vice versa. Since they deal directly with PHI in a transactional context, they are considered covered entities. Health plans, which provide or pay the cost of medical care, such as health insurance companies and HMOs, are also covered entities because they handle PHI in managing their coverage transactions.

Healthcare providers who transmit any health information in connection with transactions for which the Department of Health and Human Services has adopted standards (like billing and fund transfers electronically) are also covered entities. These transactions typically require the electronic exchange of PHI, bringing such providers under the purview of HIPAA.

However, a free health clinic may not be a covered entity if it does not engage in any covered transactions, such as billing to insurance. Many free clinics operate on a model where services are provided without charge and without billing insurance providers. Since no standard electronic transactions involving billing or claims are conducted, these clinics might not meet the definition of a covered entity under HIPAA.

It is crucial to note, however, that whether a free health clinic is a covered entity can vary. If a free clinic starts billing health insurance providers or engages in any standard electronic transactions involving PHI, it would then be considered a covered entity under HIPAA. Thus, the designation of a free health clinic as a non-covered entity is conditional and based on its operational and transactional practices.

Question: 7

Megan is believed to have alteration in sensory perception. Before the planning process begins, a major symptom must be assessed, indicating the need for a care plan. The case management team is investigating this to ensure the planning includes treating a major symptom related to this disorder. The team found:

- A major symptom of alteration in sensory perception is present

Given the information, which of the following is a major symptom the patient is likely experiencing that will be focused on in the planning process?

- A. Fear of change.
- B. A negative change in the pattern of incoming stimuli.
- C. A neutral change in the pattern of incoming stimuli.
- D. A positive change in the pattern of incoming stimuli.

Answer: B

Explanation:

Based on the information provided, the major symptom of alteration in sensory perception that Megan is experiencing, and which will be focused on in the planning process, is: *A negative change in the pattern of incoming stimuli.*

Alteration in sensory perception refers to a condition where there is a change in how a person perceives their environment. This can affect any of the senses, including vision, hearing, touch, taste, and smell. The presence of a major symptom such as a negative change in the pattern of incoming stimuli indicates a significant disruption in the normal processing of sensory information. This disruption can lead to difficulties in interpreting and responding to the environment, which can significantly impact daily functioning and quality of life.

In the context of care planning, addressing this major symptom is crucial. A negative change in sensory perception might manifest as reduced ability to detect sensory inputs, misinterpretation of those inputs, or an inability to filter out unnecessary sensory data, which can lead to sensory overload. This could result in confusion, distress, and even physical symptoms such as dizziness or nausea.

Minor symptoms that might accompany this major symptom include apathy, which refers to a lack of interest or enthusiasm; fear and anxiety, which are emotional responses to perceived threats; and disorientation, which is confusion about time, place, or identity. While these are also important to address, the primary focus of the care plan would be on mitigating the negative change in the pattern of incoming stimuli to help stabilize the patient's sensory perception.

By focusing on this major symptom, healthcare providers aim to improve Megan's ability to accurately perceive and interact with her environment, which is essential for her safety and overall well-being. This can involve various therapeutic strategies, including sensory integration techniques, environmental modifications, and possibly medication to manage any underlying causes or associated symptoms.

Question: 8

Kelly was brought into the ER with the flu. The nurse agreed she had the flu, but also felt more was wrong. The nurse suspected emotional abuse and notified the case manager. Which of the following is not an emotional abuse symptom?

- A. Act more angry than normal.
- B. Large bruises on the arm.
- C. They do not react to pain normally.
- D. They hurt themselves on purpose.

Answer: B

Explanation:

The question posed is about identifying which symptom is not indicative of emotional abuse. To address this, it's important to understand the different types of child abuse and their associated symptoms.

Child abuse is generally categorized into three main types: emotional, sexual, and physical. Each type has distinct symptoms that help in identifying the nature of the abuse: - **Emotional abuse** symptoms can be more subtle compared to physical and sexual abuse. They often manifest as changes in behavior or emotions, such as apathy (not caring much about anything), academic struggles (not doing well in school), increased anger (acting more angry than normal), abnormal reactions to pain (not reacting normally to pain), and self-harm (hurting themselves on purpose). - **Physical abuse** involves physical harm or injury to the child. Symptoms include injuries that have specific patterns (like marks from a belt or hand), injuries in usually protected areas (e.g., the back, thighs), recurrent injuries indicating ongoing abuse, and signs that the child has not received needed medical care for their injuries. - **Sexual abuse** symptoms include behavioral changes and physical signs such as reluctance to go to the bathroom, discomfort or pain when sitting, unusual discharge or bleeding from the genitals, and explicit sexual behavior or knowledge inappropriate for the child's age.

Given these categorizations, when we look back at the options provided in the question - " Act more angry than normal" is a symptom of emotional abuse. - "They do not react to pain normally" is also a symptom of emotional abuse. - "They hurt themselves on purpose" is another symptom of emotional abuse.

However, "Large bruises on the arm" does not fit within the symptoms of emotional abuse. Instead, this is a typical symptom of physical abuse. Large bruises, especially those that are patterned or in unusual places, suggest physical harm rather than emotional or psychological harm. Therefore, the correct answer to the question "Which of the following is not an emotional abuse symptom?" is "Large bruises on the arm." This symptom is indicative of physical abuse, not emotional abuse.

Question: 9

Of the following, which is the fourth most important need according to Maslow?

- A. Esteem.
- B. Self actualization.
- C. Safety.
- D. Belonging and love.

Answer: A

Explanation:

The correct answer to the question regarding the fourth most important need according to Maslow's Hierarchy of Needs is "Esteem". This hierarchy, proposed by psychologist Abraham Maslow, organizes human needs into a pyramid structure, usually with five levels. The needs are arranged in a hierarchical order from the most basic and fundamental at the bottom, to the highest level at the top. The structure of Maslow's hierarchy is as follows: 1. **Physiological Needs**: These are biological requirements for human survival, such as air, food, drink, shelter, clothing, warmth, sex, and sleep. According to Maslow, these are the foundational, most profound needs in the hierarchy and must be satisfied first. 2. **Safety Needs**: Once physiological needs are met, the second layer of human needs pertains to safety and security. This includes personal security, employment, resources, health, and property. These needs provide a sense of security and safety from physical and economic harm. 3. **Belonging and Love Needs**: After physiological and safety needs are fulfilled, the third level consists of social needs. This involves emotionally-based relationships in general, such as friendships, romantic attachments, and family. Humans need to feel a sense of belonging and acceptance among social groups, regardless if these groups are large or small. 4. **Esteem Needs**: The fourth layer includes esteem needs which are classified into two categories: (i) esteem for oneself (dignity, achievement, mastery, independence) and (ii) the desire for reputation or respect from others (e.g., status, prestige). This level of needs produces such feelings as self-worth, strength, capability, and being useful and necessary in the world. 5. **Self-Actualization Needs**: At the peak of the pyramid, self-actualization refers to the realization of a person's potential, self-fulfillment, seeking personal growth and peak experiences. It represents the growth of an individual toward fulfillment of the highest needs; those for meaning in life, in particular. Given this structure, when the question asks for the fourth most important need, the answer is "Esteem". This need pertains to the human desire for respect, self-esteem, and recognition from others, coming after the fulfillment of physiological, safety, and social belonging needs. It's crucial for building confidence and achieving personal goals, which in turn, helps pave the way to reaching the pinnacle of Maslow's hierarchy: self-actualization.

Question: 10

What does pathophysiology emphasize the most?

- A. Direct observations.
- B. Quantifiable measurements.
- C. Mechanical functions.
- D. Physical disturbances.

Answer: B

Explanation:

Pathophysiology, the scientific study of the functional changes that occur in the body as a result of a disease, places a strong emphasis on quantifiable measurements. This focus is crucial because it allows researchers and healthcare professionals to objectively evaluate how diseases alter body functions and to quantify the severity and impact of these changes.

Quantifiable measurements in pathophysiology typically involve collecting and analyzing data that can be numerically represented. This includes measurements of blood pressure, blood sugar levels, hormone levels, enzyme activities, and other biochemical markers. By using such quantifiable data, pathophysiologists can establish a baseline of what is normal and compare it against the deviations caused by diseases.

The importance of these measurements extends beyond mere diagnosis. They are essential for understanding the mechanisms underlying a disease, tracking the progression of conditions, and evaluating the efficacy of therapeutic interventions. For instance, in diabetes management, monitoring blood glucose levels helps in tailoring treatment plans and in preventing complications.

Moreover, the reliance on quantifiable measurements supports the scientific method in pathophysiology. It ensures that findings and conclusions are based on empirical evidence that can be validated and reproduced. This objectivity is vital for advancing medical knowledge and improving clinical practices.

In summary, while pathophysiology considers various factors such as mechanical functions and physical disturbances, it emphasizes quantifiable measurements the most. This emphasis facilitates a deeper understanding of diseases at a molecular and systemic level, leading to better disease management and therapeutic strategies.

Thank You for Trying Our Product

For More Information – **Visit link below:**

<https://www.examsboost.com/>

15 USD Discount Coupon Code:

G74JA8UF

FEATURES

- ✓ **90 Days Free Updates**
- ✓ **Money Back Pass Guarantee**
- ✓ **Instant Download or Email Attachment**
- ✓ **24/7 Live Chat Support**
- ✓ **PDF file could be used at any Platform**
- ✓ **50,000 Happy Customer**



Visit us at: <https://www.examsboost.com/test/acma-acm>